

Body Lift Surgery

Circumferential Body Lift | Patient Information

What is Circumferential Body Lift Surgery?

A body lift is a surgical procedure to remove excess skin and fatty tissue that may be left over after substantial weight loss. Generally, the procedure involves the removal of excess skin around the middle and lower abdomen, hips, outer thighs, back, buttocks, and the tightening of abdominal wall muscles. A body lift is not a surgical treatment for being overweight. Obese individuals who intend to lose weight should postpone all forms of body contouring surgery until they have reached a stable weight. There are a variety of different techniques used by plastic surgeons for a body lift. A circumferential body lift can be combined with other forms of body-contouring surgery, including suction-assisted lipectomy, or performed at the same time with other elective surgeries.

Is the procedure right for me?

A detailed consultation with your consultant plastic surgeon is essential prior to undergoing any surgical procedure. During the consultation, your surgeon will discuss your area/s of concern, desired improvements, options available (including no treatment at all) and expected outcomes. Your surgeon will examine you and may take some photographs for your medical records. They will ask you if you want to have someone with you during the examination and ask you to sign a consent form for taking, storing, and using the photographs. The surgeon will measure your height and weight to make sure that it is safe to do an operation. If you are overweight, or planning to become pregnant, your surgeon may suggest delaying your operation. If you decide to proceed with surgery, you will be given a 'cooling-off' period of at least 2 weeks between initial consultation and the date of your operation to ensure you have ample time to reflect on all aspects of the procedure before proceeding (although please bear in mind the waiting lists for our surgeons mean this period is invariably significantly longer).

How is the procedure carried out?

Should circumferential body lift surgery be agreed upon as the most suitable course of treatment, the procedure will be carried out by your surgeon in our state-of-the-art operating theatre. Prior to surgery, the skin is carefully marked, before anaesthetic is administered and excess skin and fat is carefully removed. Following surgery, the skin edges are closed with stitches and dressings are applied. Body lift surgery is performed as a day case under a general anaesthetic. Our highly experienced medical team will be on hand during your stay to care for your needs.

Recovery and aftercare

Following body lift surgery, patients can expect considerable bruising which will be uncomfortable and at times, painful.

Some people bruise more easily than others. The discoloration of this bruising will usually last for about a month, but the lumpiness and swelling of deep bruising can take up to 6 months to disappear. Some patients will experience temporary bruising of the genital region due to the proximity to the surgical site. It is common for the surgical site/s to ooze a little, this is usually associated with the local anaesthetic discharging but can look a little pink. Swelling at the treatment site should be expected for at least 10 days. You must expect to be left with noticeable scars. The main scar runs transversely across the lower part of the abdomen around the hips and across the top of the buttocks. Other or different scars may be left where the patient has individual problems. You can expect some numbness in the treated skin which tends to last for several months. Lipoabdominoplasty patients will have additional small scars at the site of insertion of the suction cannula. Depending on the technique used, patients should generally expect to allow for 7-10 days off work. Specific expectations on recovery time will be discussed during both initial and post-operative consultations. Light activities are comfortable in 10 to 20 days. Sports will not be possible for about 6 weeks particularly when the muscles have been strengthened with sutures. Showering is permitted on the second post-operative day. You are likely to need simple pain killers for a day or so after the operation and you will probably be asked to return a week after surgery to have sutures removed. A snug pressure garment or corset is usually advised around the lower part of the body. This is used to reduce bruising but can be taken off to wash, quickly dried and put back on. You can take this opportunity to bathe yourself. The corset is usually worn for 2 to 6 weeks.

Risks and Complications

All surgical procedures are associated with risks. Your surgeon will explain these in detail during initial consultation, enabling you to reach an informed decision on whether you wish to proceed. Serious complications are uncommon, however patients should be aware of the following complications prior to undergoing body lift surgery-

- Anaesthesia risks
- Bleeding
- Infection
- Poor scarring of skin
- Fluid accumulation (seroma)
- Pain, which may persist
- Possibility of revisional surgery
- Numbness

Other Considerations

It is important for patients to understand that body lift surgery is not a treatment for obesity. The amount of skin

T. 01904 675222

info@coppergateclinic.co.uk | www.coppergateclinic.co.uk

CC06-21



COPPERGATE
CLINIC

Body Lift Surgery

Circumferential Body Lift | Patient Information

and fat and tissue that can be removed from a localised area is limited by what is safe and there is a natural limitation when no further fat can be removed. It may not, therefore, be possible to slim down an area as much as you might like. The skin is usually tightened downwards, and this does not tighten the waist. If this is desired then one can consider removing skin vertically, but one should bear in mind that vertical scars of the abdomen are not as good. The tissue of the abdominal wall is generally fatter than the groin and if liposuction is not carried out a fatty bulge may remain above the scar. The beneficial effects of the operation will last well, however, the effects will be maintained better if the patient keeps exercising the muscles and maintains a steady weight. A further pregnancy will of course stretch the skin again, although probably not to the same degree. Patients with a BMI of more than 25 will not achieve an optimal result as the skin flaps are thicker and there is more intra-abdominal or visceral fat. Patients with a BMI of more than 30 are generally advised against having surgery until they have lost some weight.

Results

The results of all types of abdominal surgery including circumferential body lift surgery are not immediately visible as swelling can take a long time to settle, you may not see the full benefit of the operation for up to 6 months. Over time, post-surgical swelling will subside, and incision lines will fade. Satisfaction with your new image should continue to grow as you fully recover from surgery. To achieve optimal results, it is important that you follow your surgeon's post-operative instructions and follow-up visits.

Cost

Prices for body lift surgery can vary. Costs may include-

- Anaesthesia fees
- Hospital or surgical facility costs
- Medical tests
- Post-surgery garments
- Prescriptions for medication
- Surgeon's fee

After consultation and if you are planning to proceed with surgery, you will be given a written quotation regarding the cost of the planned procedure.

Alternative treatments

Depending on your specific needs and concerns, alternative surgical and non-surgical alternatives to body lift surgery may be considered. In the case of improving contour and tightening excess skin, Vaser liposuction and Renuvion skin tightening may offer a degree of benefit.

If you experience any problems or have concerns following surgery, please do not hesitate to call our direct patient line for assistance on 07908 891059.

T. 01904 675222

info@coppergateclinic.co.uk | www.coppergateclinic.co.uk

CC06-21



COPPERGATE
CLINIC

Body Lift Surgery

Circumferential Body Lift | Patient Pre & Post-Operative Care Guide

Pre-Operative Preparations

6 week period before surgery-

Patients requiring twilight or general anaesthesia should not travel long haul for the 6 week period before and 6 week period after surgery.

4 weeks before surgery-

Stop taking the contraceptive pill and HRT 4 weeks before surgery if this has been advised by your Surgeon or anaesthetist.

2 weeks before surgery- (optional)

Some patients begin taking Arnica anti-bruising tablets, three times a day, 2 weeks prior to and 2 weeks after surgery. These are available from several health and wellbeing stores, including Holland & Barratt.

1 week before surgery-

Avoid aspirin, ibuprofen, nurofen and other similar painkillers one week before and two weeks after surgery. Paracetamol is the safest painkiller to take, if necessary.

Stop taking Vitamin E and its compounds (such as evening primrose oil, fish oils, garlic and garlic capsules) 1 week before surgery and 2 weeks after surgery. Taking Vitamin E in any form can cause bleeding and post-operative complications.

1 day before and on the day of surgery-

Shower all over (including your hair and face) with Hibiscrub antiseptic liquid soap (provided on prescription from Coppergate Clinic) the day before and morning of your surgery before coming into the clinic. Do not apply moisturiser after showering.

Day of surgery-

Avoid eating solid food and milky drinks for 6 hours prior to your appointment for sedation. Water, clear drinks, black tea and black coffee can be consumed for up to 2 hours before your sedation/general anaesthetic.

Take your routine medicines at the usual time, unless advised otherwise and bring them with you (including inhalers).

Ensure at least one nail is free from acrylic or nail polish nails as a monitor will be placed on a finger.

Bring loose fitting, oversized clothing to wear after surgery.

Post-Operative Instructions

Average length of stay-

The surgery will be performed as a day case under a general anaesthetic or twilight anaesthesia unless your Surgeon advised otherwise during consultation. **If you live further than a 1 hour drive from the clinic, it is mandatory to stay in the vicinity until the following day.**

6 week period after surgery-

Patients requiring twilight or general anaesthesia should not travel long haul for the 6 week period before and 6 week period after surgery.

0-2 weeks after surgery

- It is essential to ensure activity is kept to a minimum and to rest as much as possible
- Wear the provided compression garment and TED stockings day and night, except for when showering
- Shower if you wish, but please be as quick as possible and do not linger. Dressings should be gently patted dry before the compression garment is put back on. At your follow up appointment, your surgeon will guide you on when leisurely showers and baths may resume
- Take Paracetamol as required (avoid aspirin, ibuprofen, nurofen and similar painkillers)
- Drink plenty of fluid to replace fluid removed during liposuction

0-12 weeks after surgery

- Once well-healed, apply micropore tape to scars. The tape should stay on during showers and bath and be patted dry afterwards. Tape should not be removed and replaced until it is very loose and almost falling off. Frequent removal and replacement of tape may irritate the scar.

2 weeks after surgery

- Gentle activity may resume (for example driving) but do listen to your body- it will tell you if you are overdoing things

2-4 weeks after surgery

- Wear the provided compression garment during the day

3 weeks after surgery onwards

- Moisturise and massage the umbilical scar after it is well healed

6 weeks after surgery onwards

- Static exercises such as cycling, cross training and progressive walking may resume

6-12 weeks after surgery

- Swimming and more strenuous exercise may resume, as advised by your surgeon at your follow up appointment

12 weeks after surgery onwards

- Moisturise and massage the abdominal scar and abdominal tissue

If you experience any problems or have concerns following surgery, please do not hesitate to call our direct patient line for assistance on 07908 891059.

T. 01904 675222

info@coppergateclinic.co.uk | www.coppergateclinic.co.uk

CC06-21



COPPERGATE
CLINIC